Note to Readers: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Exposure to Bisphenol A and Phthalates during Pregnancy and Ultrasound Measures of Fetal Growth in the INMA-Sabadell Cohort

Maribel Casas, Damaskini Valvi, Ana Ballesteros-Gomez, Mireia Gascon, Mariana F. Fernández, Raquel Garcia-Esteban, Carmen Iñiguez, David Martínez, Mario Murcia, Nuria Monfort, Noelia Luque, Soledad Rubio, Rosa Ventura, Jordi Sunyer, and Martine Vrijheid

Table of Contents

- Figure S1. Flowchart of the study population
- Figure S2. Fetal growth curves for femur length, head circumference, abdominal circumference, and estimated fetal weight in the INMA-Sabadell cohort
- Table S1. Variables included in each fetal growth model in the INMA-Sabadell cohort
- Table S2. Characteristics of the study population and average BPA and phthalate metabolites levels (μ g/g creatinine) by categories of the different characteristics
- Table S3. Correlation matrix of the average BPA and phthalate metabolites levels
- Table S4. Sensitivity analyses I: adjusted associations between maternal urinary BPA, MBzP, MnBP, and MiBP levels (µg/g creatinine) and fetal size and growth and birth outcomes adjusting for creatinine and excluding extreme creatinine values
- Table S5. Sensitivity analysis II: adjusted associations between maternal urinary BPA, MBzP, MnBP, and MiBP levels (μ g/g creatinine) and fetal size and growth and birth outcomes in single- and multi-pollutant models

Table S6. Adjusted associations between maternal urinary Σ DEHPm and its single metabolites (MEHP, MEHHP, MEOHP, MECPP) levels (μ g/g creatinine) and fetal size and growth in the overall population

Table S7. Adjusted associations between maternal urinary DEHPm and LMWPm levels ($\mu g/g$ creatinine) and birth outcomes in the overall population, in girls, and in boys

Table S8. Adjusted associations between maternal urinary MBzP, \sum LMWPm and its single metabolites (MEP, MiBP, MnBP) levels (μ g/g creatinine) and fetal size and growth in the overall population